## **CONTENTS**

| Special Issue: A Collection of Papers Presented at the XIVth International Carbohydrate Symposium, Stockholm, Sweden, August 14-19, 1988.  |       |
|--|-------|
| (Abstracts/contents lists published in: Chem. Abstr.; Curr. Contents; Phys. Chem. Earth Sci., Life S   | Sci.) |
| Publisher's Note   | ix    |
| Separation of fucosylated oligosaccharides using high-pH anion-exchange chromatography with pulsed-amperometric detection  M. R. HARDY AND R. R. TOWNSEND (Baltimore, MD, U.S.A.)  | 1     |
| Enzymic synthesis of di- and tri-saccharide glycosides, using glycosidases and β-D-galactoside 3-α-sialyltransferase  K. G. I. Nilsson (Arlöv, Sweden)   | 9     |
| Synthesis of pseudo-C-nucleosides  A. P. RAUTER (Lisbon, Portugal), J. A. FIGUEIREDO, AND I. M. ISMAEL (Covilhã, Portugal)   | 19    |
| Oxovanadium(IV) and copper(II) coordination by D-galacturonic and D-glucuronic acid G. MICERA, A. DESSI (Sassari, Italy), H. KOZLOWSKI, B. RADOMSKA, J. URBANSKA (Wrocław, Poland), P. DECOCK, B. DUBOIS, AND I. OLIVIER (Villeneuve d'Ascq, France)                 | 25    |
| Regioselective benzoylations of glycopyranosylamines: synthesis of partially protected glycopyranosyl isothiocyanates  J. Fuentes Mota, J. M. Garcia Fernandez, C. Ortiz Mellet, M. A. Pradera Adrian (Seville, Spain), and R. Babiano Caballero (Badajoz, Spain)    | 35    |
| Irregularity of interaction between homologous ω-aminated 1-methoxyalkyl β-D-glucopyranosides and β-D-glucosidase from sweet almond emulsin  J. LEHMANN AND L. ZISER (Freiburg, F.R.G.)  | 45    |
| Synthesis of β-nitropyrroles from amino sugars and nitroacetone JL. Chiara, A. Gómez-Sánchez, FJ. Hidalgo, and I. Yruela (Seville, Spain)  | 55    |
| Synthesis of 3-, 4-, and 6-monodeoxysucroses G. DESCOTES, J. MENTECH, AND N. ROQUES (Villeurbanne, France)   | 63    |
| A facile, regio- and stereo-selective synthesis of ganglioside GM <sub>3</sub> T. MURASE, H. ISHIDA, M. KISO, AND A. HASEGAWA (Gifu, Japan)  | 71    |
| Preparation of pyranoid glycal derivatives from phenyl thioglycosides and glycosyl phenyl sulphones  |       |
| A. FERNANDEZ-MAYORALAS, A. MARRA, M. TRUMTEL, A. VEYRIÈRES, AND P. SINAŸ (Paris, France)   | 81    |
| Isolation and identification of 3-deoxy-D-threo-hexulosonic acid as a constituent of the lipopolysac-<br>charide of Vibrio parahaemolyticus serotypes O7 and O12<br>S. Kondo, U. Zähringer, E. Th. Rietschel (Borstel, F.R.G.), and K. Hisatsune<br>(Saitama, Japan) | 97    |
| Lipopolysaccharides of <i>Proteus penneri</i> species novum  K. ZYCH AND Z. SIDORCZYK (Łódź, Poland)   | 105   |
| Selective fragmentation of methylated tobacco extracellular arabinoglucuronomannan by cleavage of β-D-glucopyranosiduronic acid units G. O. ASPINALL, L. KHONDO, AND V. PUVANESARAJAH (Toronto, Ontario, Canada) .   | 113   |
| On the specificity of the binding of cations to carrageenans: counterion n.m.r. spectroscopy in mixed carrageenan systems  L. PICULELL, S. NILSSON, AND P. STRÖM (Lund, Sweden)  | 121   |
| (Continued event   |       |

## (contents continued)

| An arabinogalactan-protein from the pulp of grape berries  L. SAULNIER AND JM. BRILLOUET (Montpellier, France)   | 137 |
|--|-----|
| Structure of the capsular polysaccharide of <i>Klebsiella ozaenae</i> serotype K4 containing 3-deoxy-D-glycero-D-galacto-nonulosonic acid Yu. A. Knirel, N. A. Kocharova, A. S. Shashkov, N. K. Kochetkov (Moscow, U.S.S.R.), V. A. Mamontova, and T. F. Solov'eva (Vladivostok, U.S.S.R.) | 145 |
| Investigation of the structure of the capsular polysaccharide of <i>Escherichia coli</i> K55 using <i>Klebsiella</i> bacteriophage φ5  A. N. Anderson and H. Parolis (Grahamstown, South Africa)   | 157 |
| Computer-assisted structural analysis of polysaccharides with an extended version of CASPER using <sup>1</sup> H- and <sup>13</sup> C-n.m.r. data PE. Jansson, L. Kenne, and G. Widmalm (Stockholm, Sweden)  | 169 |
| Preparation of oligosaccharides with β-D-mannopyranosyl and 2-azido-2-deoxy-β-D-mannopyranosyl residues by inversion at C-2 after coupling S. DAVID, A. MALLERON, AND C. DINI (Orsay, France)  | 193 |
| Notes  |     |
| An inexpensive route to 2-azido-2-deoxy-D-mannose and its conversion into an azido analog of N-acetylneuraminic acid C. Augé, S. David, and A. Malleron (Orsay, France)  | 201 |
| Comparisons of rigid and relaxed conformational maps for cellobiose and maltose A. French (New Orleans, LA, U.S.A.)  | 206 |
| Isolation and identification of 3-deoxy-7-O-α-D-galactopyranosyl-D-manno-2-octulopyranosonate from the inner core region of the lipopolysaccharide of Escherichia coli EH100 O. Holst, E. Röhrscheidt-Andrzejewski, HP. Cordes, and H. Brade (Borstel, F.R.G.)                             | 212 |
| Methylation analysis of non-glycosidic phosphoric diesters of carbohydrates O. Holst, H. Moll, H. Brade (Borstel, F.R.G.) and J. Thiem (Münster, F.R.G.)   | 219 |
| Stereoselective hydrolysis of radiolabelled <i>O</i> -pivaloyl derivatives of methyl 2-acetamido-2-deoxy-D-glucopyranoside S. TOMIĆ, Đ. LJEVAKOVIĆ, AND J. TOMŠIĆ (Zagreb, Yugoslavia)   | 222 |
| Improved synthesis of the 2-, 3-, and 4-deoxy derivatives from methyl β-D-galactopyranoside T. H. Lin, P. Kováč, and C. P. J. Glaudemans (Bethesda, MD, U.S.A.)  | 228 |
| N.m.r. studies of composition and side-chain arrangement in Sesbania aegyptiaca seed galacto-mannan  |     |
| A. K. GUPTA (West Lafayette, IN, U.S.A.) and H. GRASDALEN (Trondheim, Norway).   | 239 |
| Affinity labelling of Schizophyllum commune cellulase with [1-3H]-4,5-epoxypentyl β-cellobioside: synthesis of inhibitor and stoichiometry of interaction  A. J. CLARKE AND H. STRATING (Guelph, Ontario, Canada)  | 245 |
| Author index   | c1  |
| Subject index  | c2  |



